



牙齿/口腔健康问题

猫犬口臭

尽管口臭可因其他疾病而发生，如慢性肾病，但最常起源于口腔。¹ 当食物残渣滞留在牙齿中，或老年宠物唾液分泌减少时，可能会产生恶臭。¹ 然而，更严重的口臭通常与牙周病的发生和进展有关。牙菌斑细菌主要是厌氧菌，随着牙菌斑继续在牙齿上积聚而更加占据主导地位，它们在口腔中代谢蛋白质，导致释放出挥发性硫化物（如硫化氢），这正是产生难闻气味的原因。¹ 口臭通常是主人最先观察到的宠物牙周病症状。^{2,3} 作为综合性牙科卫生计划的一部分，营养可以在宠物口臭以及相关牙周病的管理中发挥重要作用。



重要信息

- 研究表明，投喂专门配制的磨牙咀嚼物和饮食可减少牙周病引起的口臭。^{2,4-7} 借助特定的成分和机械性能，磨牙咀嚼物和饮食可抑制导致口臭的挥发性硫化物的产生：
 - 可溶性锌盐（如硫酸锌）与硫结合，不会形成挥发性硫化物，而是形成没有气味的化合物。^{2,3} 由于锌盐具有抗菌活性，也可减少牙菌斑。^{2,3,7,8}
 - 无机磷酸盐，例如焦磷酸钠，可抑制牙菌斑矿化形成牙石。^{2,3,9} 牙石成为了滋生更多牙菌斑的病灶，促进释放更高浓度的挥发性硫化物，并导致牙周病进一步恶化。^{3,10}
 - 由于延长了咀嚼时间并增加了与牙齿和牙龈的接触，专门配制的磨牙饮食和咀嚼物可提供研磨（机械）作用，在宠物咀嚼时刮去牙菌斑和牙垢。^{2,4,7-9}
 - 研究表明磨牙咀嚼物和饮食可减少猫的牙菌斑和牙石积聚，^{11,12} 从而减轻口臭，但此益处的相关支持数据尚未发表。

(接下页)

参考文献

1. Eubanks, D. L. (2006). Canine oral malodor. *Journal of the American Animal Hospital Association*, 42, 77–79.
2. Mateo, A., Torre, C., Crusafont, J., Sallas, A., & Jeusette, I. C. (2020). Evaluation of efficacy of a dental chew to reduce gingivitis, dental plaque, calculus, and halitosis in toy breed dogs. *Journal of Veterinary Dentistry*, 37(1), 22–28. doi: 10.1177/0898756420926766
3. Roudebush, P., Logan, E., & Hale, F. A. (2005). Evidence-based veterinary dentistry: A systematic review of homecare for prevention of periodontal disease in dogs and cats. *Journal of Veterinary Dentistry*, 22(1), 6–15.
4. Gorrel, C., Warrick, J., & Bierer, T. L. (1999). Effect of a new dental hygiene chew on periodontal health in dogs. *Journal of Veterinary Dentistry*, 16(2), 77–81.
5. Simone, A., Jensen, L., Setser, C., Smith, M., & Suelzer, M. (1994) Assessment of oral malodor in dogs. *Journal of Veterinary Dentistry*, 11(2), 71–74.
6. Gorrel, C., & Bierer, T. L. (1999). Long term effects of a dental hygiene chew on the periodontal health of dogs. *Journal of Veterinary Dentistry*, 16(3), 109–113.
7. Jeusette, I. C., Román, A. M., Torre, C., Crusafont, J., Sánchez, N., Sánchez, M. C., Pérez-Salcedo, L., & Herrera, D. (2016). 24-hour evaluation of dental plaque bacteria and halitosis after consumption of a single placebo or dental treat by dogs. *American Journal of Veterinary Research*, 77(6), 613–619. doi: 10.2460/ajvr.77.6.613
8. Logan, E. I. (2006). Dietary influences on periodontal health in dogs and cats. *Veterinary Clinics of North America: Small Animal Practice*, 36, 1385–1401. doi: 10.1016/j.cvsm.2006.09.002
9. Hennet, P., Servet, E., Soulard, Y., & Biourge, V. (2007). Effect of pellet food size and polyphosphates in preventing calculus accumulation in dogs. *Journal of Veterinary Dentistry*, 24(4), 236–239. doi: 10.1177/089875640702400405
10. Harvey, C. E. (1998). Periodontal disease in dogs: Etiopathogenesis, prevalence, and significance. *Veterinary Clinics of North America: Small Animal Practice*, 28(5), 1111–1128.
11. Ingham, K. E., Gorrel, C., & Bierer, T. L. (2002). Effect of a dental chew on dental substrates and gingivitis in cats. *Journal of Veterinary Dentistry*, 19(4), 201–204.
12. Clarke, D. E., Servet, E., Hendriks, W., Thomas, D. G., Weidgraaf, K., & Biourge, V. C. (2010). Effect of kibble size, shape, and additives on plaque in cats. *Journal of Veterinary Dentistry*, 27(2), 84–89.

Purina Institute 提供易于掌握的科学信息, 帮助宠物活得更长寿、更健康, 促进人们在讨论宠物健康时将营养放在第一位。